

Management of diabetic ketoacidosis - effect of a quality improvement programme and its long-term follow-up

Kempegowda, Punith; Coombs, Benjamin; Chandan, Joht Singh; Al-Sheikhli, Jaffar; Theivendran, Kasun; Shyamanur, Bhavana; Ghosh, Sandip; Narendran, Parth

DOI:

[10.7861/clinmedicine.17-3-s8](https://doi.org/10.7861/clinmedicine.17-3-s8)

License:

None: All rights reserved

Document Version

Publisher's PDF, also known as Version of record

Citation for published version (Harvard):

Kempegowda, P, Coombs, B, Chandan, JS, Al-Sheikhli, J, Theivendran, K, Shyamanur, B, Ghosh, S & Narendran, P 2017, 'Management of diabetic ketoacidosis - effect of a quality improvement programme and its long-term follow-up', *Clinical Medicine*, vol. 17, no. Suppl 3, s8. <https://doi.org/10.7861/clinmedicine.17-3-s8>

[Link to publication on Research at Birmingham portal](#)

Publisher Rights Statement:

Checked for eligibility: 15/04/2019

General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact UBIRA@lists.bham.ac.uk providing details and we will remove access to the work immediately and investigate.

Management of diabetic ketoacidosis – effect of a quality improvement programme and its long-term follow-up

Authors: Punith Kempegowda,^{1,2} Benjamin Coombs,¹ Joht Singh Chandan,^{1,2} Jaffar Al-Sheikhli,¹ Kasun Theivendran,¹ Bhavana Shyamanur,¹ Sandip Ghosh¹ and Parth Narendran^{1,2}

Aims

To study the effect of a quality improvement programme (QIP) on the overall care for patients admitted with diabetic ketoacidosis (DKA).

Methods

Patients aged ≥ 17 years admitted with DKA from April 2014 to September 2016 at a large teaching hospital in the West Midlands were included in the study. Patients managed in intensive care units were excluded to avoid bias from one-to-one care. Various interventions were introduced from October 2014 to March 2015 to improve DKA care as part of the QIP. The study period was divided into five groups of 6 months: baseline (52), intervention (52), early (53), intermediate (66) and late- (36) follow-up period. Data measuring socio-demographic profile (age, gender, ethnicity) and various aspects of DKA management (presenting aetiology, glucose and ketone monitoring, prescribing fixed-rate insulin infusion and fluids) were collected and analysed using Microsoft Excel (version 2010) Statistical Analysis Software Package version 23 (IBM Corp., USA). The primary outcome of the study was measured by the effect on DKA duration. Results are expressed in percentage and median with interquartile range (IQR) as appropriate.

Results

A total of 259 DKA episodes (male:female, 1:1.14) were analysed. There was no significant difference in the age between the groups. Poor compliance (32.1%) was the most common precipitating factor of DKA during the study. Glucose was monitored hourly in 91.7% (IQR, 83.3–108.2%) in pre-intervention, 91.7% (IQR, 75.0–114.6%) in intervention, 116.7% (IQR, 91.7–150.5%) of early, 125.2% (IQR, 100.1–149.1%) in intermediate and 120.3% (IQR, 91.4–143.9%) in late follow-up groups respectively. Hourly ketone monitoring was met in 58.3% (IQR, 25.0–80.6%) in pre-intervention, 50.0% (IQR,

33.3–73.0%) in intervention, 42.2% (IQR, 26.2–66.9%) in early, 61.3% (IQR, 41.2–83.8%) in intermediate and 51.5% (IQR, 26.3–71.4%) in late follow-up patients respectively.

The duration of DKA during the five groups as defined above were 20.5 hours (IQR, 12.5–41.1 hours), 16.6 hours (IQR, 11.9–27.9 hours), 11.5 hours (IQR, 7.1–20.7 hours), 7.4 hours (IQR, 5.0–11.3 hours) and 9.0 hours (IQR, 6.8–15.6 hours) respectively. Overall, there was statistically significant reduction in DKA duration following intervention, and the effect of QIP persisted up to 1 year post-intervention.

Conclusions

A well-designed QIP based on specific targets and continued feedback improved the quality of care for DKA. Although there was a significant reduction in DKA duration through our QIP, an increasing DKA duration trend during long-term follow-up suggests the need for ongoing interventions to maintain the improvements. ■

Conflict of interest statement

None.

Authors: ¹University Hospitals Birmingham NHS Foundation Trust, Birmingham, UK; ²Institute of Metabolism and Systems Research, University of Birmingham, Birmingham, UK